

Infection Control Policy

Contents:

Statement of intent

1. Legal framework

Preventative measures

2. Ensuring a clean environment
3. Child immunisation
4. Staff immunisation
5. Contact with animals
6. Water-based activities

In the event of infection

7. Preventing the spread of infection
8. Vulnerable children
9. Procedures for unwell children, staff and beneficiaries
10. Exclusion
11. Medication
12. Outbreaks of infectious diseases
13. Pregnant staff members
14. Staff handling food
15. Managing specific infectious diseases
16. Monitoring and review

Appendices

- A. Managing specific infectious diseases
- B. Infection absence periods
- C. Diarrhoea and vomiting outbreak action checklist

Statement of intent

Infections can easily spread within the charity due to:

- Children's undeveloped immune systems.
- The close-contact nature of the childcare and parent carer & toddler group environment.



- Some children having not yet received full vaccinations.
- Poor understanding of good hygiene practices.

Infections commonly spread in the following ways:

- **Respiratory spread** – contact with coughs or other secretions from an infected person.
- **Direct contact spread** – direct contact with the infecting organism, e.g. skin-on-skin contact during sports.
- **Gastrointestinal spread** – contact with contaminated food or water, or contact with infected faeces or unwashed hands.
- **Blood-borne virus spread** – contact with infected blood or bodily fluids, e.g. via bites or used needles.

The charity actively prevents the spread of infection via the following measures:

- Maintaining high standards of personal hygiene and practice
- Maintaining a clean environment
- Encouraging routine immunisations
- Taking appropriate action when infection occurs

This policy aims to help staff prevent and manage infections in the charity. It is not intended to be used as a tool for diagnosing disease, but rather a series of procedures informing staff what steps to take to prevent infection and what actions to take when infection occurs.

1. Legal framework

This policy has due regard to legislation including, but not limited to, the following:

- The Control of Substances Hazardous to Health Regulations (COSHH) 2002 (amended 2004)
- Health and Safety at Work etc. Act 1974
- The Management of Health and Safety at Work Regulations 1999
- The Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (RIDDOR) 2013
- The Health Protection (Notification) Regulations 2010
- The Health Protection (Local Authority Powers) Regulations 2010

This policy has due regard to statutory guidance including, but not limited to, the following:

- UK Health and Security Agency (2023) 'Health protection in children and young people settings, including education'

This policy operates in conjunction with the following charity policies and documents:

- Health and Safety Policy



- Administering Medication Policy
- First Aid Policy
- Swimming Risk Assessment
- Bodily Fluids Risk Assessment

Preventative Measures

2. Ensuring a clean environment

Handwashing

The charity will ensure everyone on the premises will have access to liquid soap, warm water and paper towels. Staff will check, encourage and supervise handwashing in children where appropriate.

All staff and children will be advised to wash their hands after using the toilet, before eating or handling food, after playing in the garden and after touching animals.

Cleaning

All cleaning staff will be appropriately trained and appropriate PPE, e.g. gloves, aprons and surgical masks, will be available. The line manager will devise a cleaning schedule that clearly describes the activities required, the frequency of cleaning and who will carry out which activities. Cleaning standards will be regularly monitored to ensure effectiveness and that all areas or surfaces in contact with food, dirt or bodily fluids are regularly cleaned and disinfected.

Cleaning equipment will be maintained to a high standard and is colour coded according to area of use. Line managers will be responsible for monitoring cleaning standards and discussing any issues that may arise with staff members.

Sanitary facilities

Wall-mounted soap dispensers or free standing bottles of liquid soap will be used in all toilets – bar soap is never used.

A foot-operated waste-paper bin will be available where disposable paper towels are used.

Toilet paper will always be available in cubicles. Suitable sanitary disposal facilities will be provided where necessary, including where there are children aged nine and above.

Nappy changing areas for childcare

A designated changing area will be established away from play facilities and food and drink areas, and with appropriate hand washing facilities.



Children's skin will be cleaned with disposable wipes, and nappy creams and lotions will be labelled with the relevant child's name and not shared with others.

Changing mats will be wiped with soapy water or a mild detergent wipe after each use. If a mat is visibly soiled, it will be cleaned thoroughly with hot soapy water immediately. Mats will be checked on a weekly basis for tears and damage and replaced if necessary.

Hand wash basins are not used for cleaning potties, and instead staff will use a designated sink located in the area where potties are used. Potties will be washed in hot, soapy water, dried and stored upside down. When cleaning potties, rubber gloves will be used to flush waste down the toilet. Rubber gloves will be washed after use whilst still being worn and dried, and the wearer will wash their hands afterwards.

Soiled nappies will be disposed of in the nappy waste bins inside a wrapped plastic bag.

Continence aid facilities

Children who use continence aids, e.g. continence pads and catheters, will be encouraged to be as independent as possible. Pads will be changed in a designated area with adequate handwashing facilities, and disposable powder-free latex gloves and a disposable plastic apron will be worn.

Laundry

All laundry will be washed in a separate dedicated facility, and any soiled linens will be washed separately.

Manual sluicing of clothing will not be permitted, and gloves and aprons will be worn when handling soiled linen or clothing. Hands will be thoroughly washed after gloves are removed.

Ventilation

Indoor spaces will be kept well-ventilated to help reduce the amount of respiratory germs. Areas of the charity where there may be poor ventilation will be identified, e.g. through the use of CO2 monitors, and appropriate action taken, e.g. partially opening windows and doors to let fresh air in. The need for increased ventilation will always be balanced against the need to maintain a comfortable temperature for staff, children, beneficiaries and visitors.

Toys and other equipment

When purchasing toys, the charity will ensure they all carry a BS, BSI or CE mark and that, where possible, they can be easily cleaned. Toys will be stored in clean containers. Children will not be allowed to take them into toilet areas.



A written schedule will be put in place to ensure that toys and equipment are cleaned on a daily basis with a deep clean termly. Toys that are “soft”, e.g. modelling clay, will be discarded whenever they look dirty.

Sandpits will be covered when not in use and the sand is changed termly for indoor sandpits and, for outdoor sandpits, as soon as the sand becomes discoloured or malodorous. Sand will be sieved or raked on a weekly basis.

Water play troughs will be emptied, washed with detergent and hot water, dried and stored upside-down when not in use for long periods. When in use, the water will be replenished, at a minimum, on a daily basis, and the trough will remain covered overnight.

Managing cuts, bites, and bodily fluid spills

Standard precautions will always be taken when dealing with any cuts and abrasions.

Any spillages of blood, faeces, saliva, vomit, or nasal discharges will be cleaned immediately in line with the Bodily Fluids Risk Assessment.

PPE will be worn where there is a risk of contamination with blood or bodily fluids. Gloves are disposable, non-powdered vinyl or latex and CE marked. If there is a risk of splashing to the face, disposable eye protection, or reusable eye protection that is decontaminated prior to next use, will be worn.

Cuts and abrasions will be cleaned under running water or using a disposable container with water and wipes. The wound will be carefully dabbed dry then covered with a waterproof dressing or plaster. The dressing will be changed as often as is necessary. Staff will wear disposable gloves when in contact with any accident or injury, e.g. washing grazes, or dressing wounds.

If a staff member, child or beneficiary suffers a bite or scratch that does not break the skin, the affected area will be cleaned with soap and warm running water. If a bite, scratch or puncture injury breaks the skin or may have introduced someone else’s blood, the affected area will be washed thoroughly with soap and warm running water, the incident will be recorded in the accident book or accident log in childcare, the wound will be covered with a waterproof dressing, and medical advice sought immediately.

When coughing or sneezing, all staff, children and beneficiaries will be encouraged to cover their nose and mouth with a disposable tissue and dispose of the tissue after use, and to wash their hands afterwards.

Safe management of waste – including sharps

The charity will ensure that all waste produced is dealt with by a licensed waste management company.



Any PPE used will be placed in a refuse bag and disposed of as normal domestic waste. PPE should not be put in a recycling bin or dropped as litter.

Injuries incurred through sharps found on the charity premises will be treated in line with the charity's Health and Safety Policy. All sharps found on the premises will be disposed of in the sharps bin whilst wearing PPE.

3. Child immunisation

The charity is aware of the vital role it plays in supporting the routine immunisation programme and will liaise with local health services to share information with parents at key points.

Early Years Foundation Stage (EYFS) children's immunisation status will be checked upon registration.

The charity including childcare staff and Family Support staff will keep up-to-date with national and local immunisation scheduling and advice via www.nhs.uk/conditions/vaccinations/. Below is a list of vaccines available on the NHS, including who should have them and when:

- **EYFS Children** - Before starting school, children should be given their second injection of the MMR vaccine, usually at 3 years and 4 months. Children should also be given their 4-in-1 pre-school booster against diphtheria, tetanus, whooping cough and polio, usually at 3 years and 4 months.
- **Primary school aged children** - All children in Reception to Year 4 will be offered nasal flu vaccinations annually.
- **Secondary school aged children** - Children aged between 12 and 13 can choose to get the HPV vaccine to protect themselves against cervical cancer, some mouth and throat cancers and some cancers of the anal and genital areas. This vaccine comprises two injections given 6-12 months apart.
- **Secondary school aged children** - All children aged 14 will be offered the 3-in-1 teenage booster vaccination to top-up the effects of the pre-school vaccines against diphtheria, polio and tetanus.
- **Secondary school aged children** - All children aged 14 will be offered the MenACWY vaccine as part of the routine adolescent schools programme.

The charity will work with local health professionals to signpost parents to where they can book coronavirus (COVID-19) vaccination appointments for children online, at a vaccination centre or pharmacy, or at a local walk-in centre without an appointment.

The charity will notify the local DfE team or Early Years team of any anti-vaccination activity, e.g. campaign letters and emails spreading misinformation about vaccination programmes.



Only information from trusted sources, e.g. the NHS, and where its authenticity is assured will be shared by the charity.

4. Staff immunisation

Staff will be encouraged to check their immunisation records and contact their GP practice if they are unsure if they are up to date with vaccinations or if they need to catch up.

MMR vaccination: In particular, all staff should make sure that they have had 2 doses of the MMR vaccine. The MMR vaccine is the safest and most effective way to protect against measles, mumps, and rubella, which are viral infections that can quickly spread and cause outbreaks. Rubella caught in pregnancy can lead to miscarriage or cause very serious harm to the unborn baby and so all people should make sure they are up to date with their 2 doses, staff will be encouraged to check their immunisation status and contact their GP for any further medical advice.

Hepatitis B: The hepatitis B vaccine is recommended for staff who are involved in the care of individuals with severe learning disability or challenging behaviour who live in institutional accommodation. In all circumstances the charity will undertake an occupational health risk assessment and pay for the vaccine if it is required.

Where necessary, staff will be permitted time off to receive any advised immunisations.

5. Contact with animals

Animals in the charity will be strictly controlled under our Animals in the charity risk assessment.

The line manager will assign a member of staff with suitable knowledge and experience to be responsible for any onsite animals and abide by the Animal Welfare Act 2006, which places a duty on animal owners to ensure their animal's welfare needs are met.

Where a staff member, child or beneficiary has a support animal, a member of staff will be assigned responsibility for infection prevention measures and supporting them to follow these.

The charity will ensure it has the correct insurance arrangements in place for any animals brought onto the premises.

Only mature and toilet trained animals will be allowed on the premises.

Animals will always be supervised when in contact with children and anyone handling animals will wash their hands immediately after touching them, their bedding or equipment.

Pregnant staff will be advised to avoid contact with any animal on the premises due to the risk of toxoplasmosis.



Visits to farms and zoos will be suitably risk assessed.

6. Water based activities

Swimming activities

General swimming activities will be governed by the control measures outlined in a Swimming Risk Assessment.

Anyone who has experienced vomiting or diarrhoea preceding the swimming activity will not be permitted to attend public swimming pools until two weeks after the end of symptoms.

Other activities

Alternative water-based activities will only be undertaken at reputable centres.

Attendees will cover all cuts, scratches and abrasions with waterproof dressings before taking part and wash their hands immediately after the activity. No food or drink will be consumed until hands have been washed.

After canoeing or rowing, attendees will immediately wash or shower.

If a member of the charity becomes ill within three to four weeks of an activity taking place, the charity will encourage them or in the case of a child their parents to seek medical advice and inform the treating doctor of theirs or their child's participation in these activities.

In the event of infection

7. Preventing the spread of infection in children

Parents will be asked not to bring their child into the setting in the following circumstances:

- The child shows signs of being unwell and needing one-to-one care.
- The child has taken, or needs to take, infant paracetamol, ibuprofen or 'Calpol'.
- The child has a high temperature or fever.
- The child has been vomiting and/or had diarrhoea within the last 48 hours.
- The child has an infection and the recommended exclusion period stated in the 'Appendix A - Managing specific infectious diseases' section has not yet passed.

8. Vulnerable adults and children

Vulnerable adults and children with impaired immune defence mechanisms, known as immunosuppressed, are more likely to acquire infections. In addition, the effect of an infection is likely to be more significant for such vulnerable adults and children. These may have a disease that compromises their immune system or be undergoing treatment, e.g. chemotherapy, that has a similar effect.



The charity will be notified if someone is “vulnerable”. Parents are responsible for notifying the charity if their child is vulnerable.

If a vulnerable adult is thought to have been exposed to an infectious disease, they will be encouraged to seek medical advice from their doctor or specialist. If a vulnerable child is thought to have been exposed to an infectious disease, the child’s parents will be informed and encouraged to seek medical advice from their doctor or specialist.

9. Procedures for unwell staff, children and beneficiaries

Staff will be required to know the warning signs of children becoming unwell including, but not limited to, the following:

- Not being themselves.
- Not eating, e.g. at snack or lunchtimes.
- Wanting more attention or sleep than usual.
- Displaying physical signs of being unwell, e.g. watery eyes, a flushed face or clammy skin.

Where a staff member identifies someone in the setting as unwell, their temperature will be taken by the staff member trained in first aid, if it is a child and their parents are not present, they will be informed of the situation immediately by reception staff.

The staff member trained in first aid will also:

- Attempt to cool the person down if they are too hot, by opening a window and suggesting that they remove their top layers of clothing such as coats, jumpers.
- Provide the person with a drink of water.
- Move the person to a quieter area of the room or building.
- In the case of a child, ensure there is a staff member available to comfort the child.
- Summon emergency medical help if required.

Children, staff and beneficiaries displaying any of the signs of becoming unwell outlined above will be sent home, and the charity will recommend that they see a doctor.

If a child is identified with sickness and diarrhoea, the child’s parents will be contacted immediately and the child will be sent home and may only return after 48 hours have passed without symptoms.

If a staff member is suffering from vomiting and diarrhoea, they will be sent home and may not return until 48 hours have passed without symptoms.

If a beneficiary is suffering from vomiting and diarrhoea, they will be sent home and may not return until 48 hours have passed without symptoms.



If the charity is unable to contact a child's parents in any situation, the child's alternative emergency contacts will be contacted.

Contaminated clothing

If the clothing of a staff member or a child who is sick becomes contaminated, the clothing will be removed as soon as possible and placed in a plastic bag. The clothing will be sent home with the child, and parents are advised of the best way to launder the clothing.

Contaminated clothing will be washed separately in a washing machine, using a pre-wash cycle on the hottest temperature that the clothes will tolerate.

10.Exclusion

Children, staff and beneficiaries who are showing the symptoms of an infectious disease or have been diagnosed by a health professional or diagnostic test will be advised to stay away from the charity for the minimum period recommended, if required, and until well enough.

The charity will follow the recommendations of the UKHSA in determining the minimum required period for staff, children and beneficiaries to stay away from the setting following an infection, as laid out in Appendix B - infection absence periods.

The charity will expect parents to agree that, if their child is unwell and has symptoms of an infectious illness, such as a fever, they should not attend the setting, given the potential risk to others.

If a parent insists on a child with symptoms attending the setting, where they have a confirmed or suspected case of an infectious illness, the charity will take the decision to exclude the child from the charity – on medical grounds - if, in the charity's reasonable judgement, it is necessary to protect other children, staff and other beneficiaries from possible infection.

For some infections, individuals may be advised to remain away from the setting for a longer period of time and the charity will follow any advice received from the local health protection team (HPT).

If a child, member of staff or beneficiary is a close contact of someone unwell with an infectious disease, but is not confirmed to be infected, this is not normally a valid reason for exclusion; however, the local HPT may advise on specific precautions to take in response to a case or outbreak.

The charity is aware that exclusion may cause challenges for parents due to unexpected time off and, that some children may become vulnerable to domestic abuse or neglect during times where they would usually be at the setting. When recommending exclusion on public health grounds, the charity will work with their HPT to consider any adverse effects or hidden harms



a child may be exposed to by imposing isolation, and staff will be alert and proactive in sharing information as early as possible. The charity will always follow procedures outlined in the charity's child protection policies if they have a concern about a child.

The charity is aware that exclusion on public health grounds may cause some children or staff members to feel isolated or anxious. In such situations, the charity will signpost them to mental health and wellbeing support services.

11. Medication

Where a child has been prescribed medication by a doctor, dentist, nurse or pharmacist, the first dose will be given at home, in case the child has an adverse reaction.

The child will only be allowed to return to the setting 24 hours after the first dose of medication, to allow it time to take effect.

All medicine provided in the charity will be administered in line with the Administering Medication Policy.

12. Outbreaks of infectious diseases

An incident is classed as an 'outbreak' where two or more people experiencing a similar illness are linked in time or place, or a greater than expected rate of infection is present compared with the usual background rate, such as:

- Two or more people in the same room are suffering from vomiting and diarrhoea.
- A greater number of children than usual is diagnosed with scarlet fever.
- There are two or more cases of measles at the setting.

Where an outbreak is suspected (even if it cannot be confirmed), the CEO will promptly contact the HPT to discuss the situation and agree if any actions are needed. The charity will support the HPT's identified control measures with clear and prompt communication with staff, parents and beneficiaries and rapid coordination of arrangements, e.g. staff immunisation.

The CEO will provide the following information:

- The number of staff, children and beneficiaries affected
- The symptoms present
- The date the symptoms first appeared
- The number of people affected

The HPT will provide the charity with draft letters and factsheets to distribute to staff, parents and beneficiaries.



The HPT will always treat outbreaks in the strictest confidence; therefore, information provided to staff, parents and beneficiaries during an outbreak will never include names and other personal details.

If a member of staff suspects the presence of an infectious disease in the setting, they will contact the CEO for further advice.

If a parent or beneficiary informs the charity that they or their child carries an infectious disease, other attendees will be observed for similar symptoms by staff.

If a child is identified as having a notifiable disease, as outlined in Appendix B - Infection Absence Periods, the charity will inform the parents, who should inform their child's GP. It is a statutory requirement for doctors to then notify their local UK Health Security Agency centre.

During an outbreak, enhanced or more frequent cleaning protocols may be undertaken, in line with information provided by the local HPT. The CEO with line managers will liaise with the cleaning staff to ensure these take place.

Under the Health Protection (Notification) Regulations 2010, the charity will always report instances of the following diseases to the HPT:

- Acute encephalitis
- Acute meningitis
- Acute poliomyelitis
- Acute infectious hepatitis
- Anthrax
- Botulism
- Brucellosis
- Cholera
- Diphtheria
- Enteric fever (typhoid or paratyphoid fever)
- Food poisoning
- Haemolytic uraemic syndrome (HUS)
- Infectious bloody diarrhoea
- Invasive group A streptococcal disease and scarlet fever
- Legionnaires' disease
- Leprosy
- Malaria
- Measles
- Meningococcal septicaemia
- Mumps



- Plague
- Rabies
- Rubella
- SARS
- Smallpox
- Tetanus
- Tuberculosis
- Typhus
- Viral haemorrhagic fever (VHF)
- Whooping cough
- Yellow fever

13. Pregnant staff members

If a pregnant staff member develops a rash or is in direct contact with someone who has a potentially contagious rash, the charity will strongly encourage them to speak to their GP or midwife.

Pregnant staff members will be advised to ensure they are up-to-date with the recommended vaccinations, including against coronavirus.

Chickenpox: If a pregnant staff member has not already had chickenpox or shingles, becoming infected can affect the pregnancy. If a pregnant staff member believes they have been exposed to chickenpox or shingles and have not had either infection previously, they will speak to her midwife or GP as soon as possible. If a pregnant staff member is unsure whether they are immune, the charity will encourage them to take a blood test.

Measles: If a pregnant staff member is exposed to measles, they will inform their midwife immediately.

Rubella (German measles): If a pregnant staff member is exposed to rubella, they will inform their midwife immediately.

Slapped cheek disease (Parvovirus B19): If a pregnant staff member is exposed to slapped cheek disease, they will inform their midwife promptly.

14. Staff handling food

Food handling staff suffering from transmittable diseases will be excluded from all food handling activity until advised by the local Environmental Health Officer (EHO) that they are clear to return to work. Both food handling staff are not permitted to attend work if they are suffering from diarrhoea and/or vomiting. They are not permitted to return to work until 48



hours have passed since diarrhoea and/or vomiting occurred, or until advised by the local EHO that they are allowed to return to work.

The charity will notify the local Environmental Health Department as soon as they are notified that a staff member engaged in the handling of food has become aware that they are suffering from, or likely to be carrying, an infection that may cause food poisoning.

Food handlers are required by law to inform the charity if they are suffering from any of the following:

- Typhoid fever
- Paratyphoid fever
- Other salmonella infections
- Dysentery
- Shigellosis
- Diarrhoea (where the cause of which has not been established)
- Infective jaundice
- Staphylococcal infections likely to cause food poisoning like impetigo, septic skin lesions, exposed infected wounds, boils
- E. coli VTEC infection

'Formal' exclusions will be issued where necessary, but employees are expected to provide voluntary 'off work' certificates from their GP.

15. Managing specific infectious diseases

When an infectious disease occurs in the setting, staff will follow the appropriate procedures set out in the Appendix A - Managing Specific Infectious Diseases.

16. Monitoring and review

All members of staff will be required to familiarise themselves with this policy as part of their induction programme.

The CEO will review this policy on an annual basis and will make any changes necessary, taking into account the current effectiveness of infection control and prevention.

Appendix A – Managing specific infectious diseases

Disease	Symptoms	Considerations	Exclusion period
Athlete's foot	Scaling, peeling or cracking of the skin, particularly between the toes and on soles of the feet, or blisters containing fluid. The infection may be itchy, and toenails can become discoloured, thick and crumbly.	Cases are advised to see their local pharmacy or GP for advice and treatment.	Exclusion is not necessary.
Chicken pox	Sudden onset of fever with a runny nose, cough and generalised rash. The rash then blisters and scabs over. Several blisters may develop at once, so there may be scabs in various stages of development. Blisters typically crust up and fall off naturally within one to two weeks. Some mild infections may not present symptoms.	Cases are advised to consider pharmacy remedies to alleviate symptoms and consult their GP. Immediate medical advice should be sought if abnormal symptoms develop, e.g. infected blisters, chest pain or difficulty breathing.	Chickenpox is infectious from 48 hours prior to a rash appearing, and until all blisters have crusted over, typically five to six days after the onset of a rash. Cases will be excluded from the setting for at least five days from the onset of a rash and until all blisters have dried and crusted over. It is not necessary for all the spots to have healed before the case returns to the setting.
Cold sores	The first signs of cold sores are tingling, burning or itching in the affected area. Around 24 hours after the first signs appear the area will redden and swell, resulting in a fluid-filled blister or	Cases are advised not to touch the cold sore or pick at the blisters. Sufferers of cold sores should avoid kissing people and should not share	Exclusion is not necessary.

Disease	Symptoms	Considerations	Exclusion period
	blisters. After blistering, they may form ulcers, then dry up and crust over.	food and items such as cutlery, cups, towels and facecloths.	
Conjunctivitis	The eye(s) become reddened and swollen, and there may be a sticky or watery discharge. Eyes may feel itchy and 'gritty'.	<p>Cases are encouraged to seek advice, wash their hands frequently and not to rub their eyes.</p> <p>Cases will be advised to seek advice and treatment from their local pharmacist.</p> <p>The HPT will be contacted if an outbreak occurs.</p>	<p>Exclusion is not necessary.</p> <p>In the case of an unmanageable outbreak, exclusion may become necessary, as per the HPT's advice.</p>
Cryptosporidiosis	Symptoms include abdominal pain, diarrhoea and occasionally vomiting.	Individuals in the setting will be asked to wash hands regularly. Kitchen and toilet areas will be cleaned regularly.	Cases will be excluded until 48 hours have passed since symptoms were present.
Diarrhoea and vomiting (gastroenteritis)	Symptoms include diarrhoea and/or vomiting; diarrhoea is defined as three or more liquid or semi-liquid stools in a 24-hour period.	The HPT will be contacted where there are more cases than usual.	<p>Cases will be excluded until 48 hours have passed since symptoms were present – for some infections, longer periods are required, and the HPT will advise accordingly.</p> <p>If medication is prescribed, the full course must be completed and there must be no further symptoms displayed for 48 hours</p>

Disease	Symptoms	Considerations	Exclusion period
			<p>following completion of the course before the cases may return to the setting.</p> <p>Cases will be excluded from swimming activities for two weeks following their last episode of diarrhoea.</p>
E. coli STEC	Symptoms vary but include diarrhoea which can be bloody, abdominal pain, vomiting and fever.	Cases will immediately be sent home and advised to speak to their GP.	<p>Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.</p> <p>Where the sufferer poses an increased risk, e.g. food handlers, pre-school infants, vulnerable adults & children, they will be excluded until a negative stool sample has been confirmed.</p> <p>The HPT will be consulted in all cases.</p>
Food poisoning	Symptoms normally appear within one to two days of contaminated food being consumed, although they may start at any point between a few hours and several weeks later. The main symptoms are likely to be nausea, vomiting, diarrhoea, abdominal pain and fever.	<p>Cases will be sent home.</p> <p>The HPT will be contacted where two or more cases with similar symptoms are reported.</p> <p>All outbreaks of food poisoning outbreak will be investigated.</p>	<p>Cases will be excluded until 48 hours have passed since symptoms were present.</p> <p>For some infections, longer exclusion periods may be required. The HPT will advise in such cases.</p>

Disease	Symptoms	Considerations	Exclusion period
Giardiasis	Infection can be asymptomatic, and the incubation period is between 5 and 25 days. Symptoms can include abdominal pain, bloating, fatigue and pale, loose stools.	Cases will be sent home. The HPT will be contacted where two or more cases with similar symptoms are reported.	Cases will be excluded until 48 hours have passed since symptoms were present.
Glandular fever	Symptoms include severe tiredness, aching muscles, sore throat, high fever, swollen glands in the neck and occasionally jaundice.	The sufferer may feel unwell for several months with fatigue and the setting will provide reasonable adjustments where necessary.	Exclusion is not necessary, and cases can return to the setting as soon as they feel well.
Group A Streptococcus (GAS)	Symptoms include flu-like symptoms, sore throat, rough rash, scabs and sores (impetigo), pain and swelling, severe muscle aches, nausea and vomiting.	GAS can cause a number of infections, some mild and some more serious. Milder infections can be easily treated with antibiotics and usually recover at home in a few days.	Cases will be excluded for 24 hours after starting to take antibiotics.
Hand, foot and mouth disease	Symptoms include a fever, reduced appetite and generally feeling unwell. One or two days later, a rash with blisters may develop with blisters on the inside of cheeks, gums, sides of the tongue, and hands and feet. Not all cases will have symptoms.	Where rare additional symptoms develop, e.g. high fever, headache, stiff neck, back pain or other complications, prompt medical advice should be sought.	Exclusion is not necessary, and cases can return to the setting as soon as they feel well.
Head lice	Other than the detection of live lice or nits, there are no immediate symptoms until two to three	Treatment is only necessary when live lice are seen.	Exclusion is not necessary, as headlice are not considered a health hazard.

Disease	Symptoms	Considerations	Exclusion period
	<p>weeks after infection, where itching and scratching of the scalp occurs.</p>	<p>Staff are not permitted to inspect any child's hair for head lice.</p> <p>If a staff member incidentally notices head lice in a child's hair, the childcare manager will inform the parents and advise them to treat their child's hair.</p> <p>Upon noticing, staff members are not required to send the child home; the child is permitted to stay in setting for the remainder of the day.</p> <p>When a child has been identified as having a case of head lice, a letter will be sent home to all parents notifying them that a case of head lice has been reported and asking all parents to check their children's hair.</p>	<p>In severe, ongoing cases, the LA does have the power to exclude. This use of power must be carefully considered, and exclusion should not be overused.</p>
<p>Hepatitis A</p>	<p>Infection can be asymptomatic. Symptoms can include abdominal pain, loss of appetite, nausea, fever and fatigue, followed by jaundice, dark urine and pale faeces.</p>	<p>The illness in children usually lasts one to two weeks but can last longer and be more severe in adults.</p>	<p>Cases are excluded while unwell and for seven days after the onset of jaundice (or the onset of symptoms if no jaundice presents).</p>

Disease	Symptoms	Considerations	Exclusion period
Hepatitis B	Infection can be asymptomatic. Symptoms can include general fatigue, nausea, vomiting, loss of appetite, fever and dark urine, and older cases may develop jaundice. It can cause an acute or chronic illness.	<p>The HPT will be contacted where advice is required.</p> <p>The procedures for dealing with blood and other bodily fluids will always be followed.</p> <p>The accident book will always be completed with details of injuries or adverse events related to cases.</p>	<p>Acute cases will be too ill to attend the setting, and their doctor will advise when they are fit to return.</p> <p>Chronic cases will not be excluded or have their activities restricted.</p> <p>Staff with chronic hepatitis B infections will not be excluded.</p>
Hepatitis C	Symptoms are often vague but may include loss of appetite, fatigue, nausea and abdominal pain. Less commonly, jaundice may occur.	<p>The procedures for dealing with blood and other bodily fluids will always be followed.</p> <p>The accident book will always be completed with details of injuries or adverse events related to cases.</p>	Cases will not be excluded or have their activities restricted.
Impetigo	Symptoms include sores, typically on the face and on the hands and feet. After around a week, the sores burst and leave golden brown crusts and can sometimes be painful and itchy.	<p>Towels, facecloths and eating utensils will not be shared by anyone.</p> <p>Toys and equipment will be cleaned thoroughly; non-washable soft toys will be wiped or washed with a detergent using warm water and dried thoroughly.</p>	Cases will be excluded until all sores or blisters are crusted over, or 48 hours after commencing antibiotic treatment.

Disease	Symptoms	Considerations	Exclusion period
<p>Influenza</p>	<p>Symptoms include headache, high temperature, cough, sore throat, aching muscles and joints, and fatigue. Younger cases may present different symptoms, e.g. without fever but with diarrhoea.</p>	<p>Those in risk groups will be encouraged to have the influenza vaccine.</p> <p>Anyone with flu-like symptoms will stay home until they have recovered.</p>	<p>There is no specific exclusion period; cases will remain home until they have fully recovered.</p>
<p>Invasive Group A Streptococcus (iGAS)</p>	<p>Symptoms include flu-like symptoms, sore throat, rough rash, scabs and sores (impetigo), pain and swelling, severe muscle aches, nausea and vomiting.</p>	<p>These infections are caused by the bacteria getting into parts of the body where it is not normally found, such as the lungs or bloodstream. In rare cases an iGAS infection can be fatal.</p>	<p>Inform HPT if any cases reported. The HPT will carry out a risk assessment and undertake appropriate investigations and/or actions as required.</p>
<p>Measles</p>	<p>Symptoms include a runny nose, cough, conjunctivitis, high fever and small white spots inside the cheeks. Around the third day, a rash of flat red or brown blotches may appear on the face then spread around the body.</p>	<p>All children are encouraged to have MMR immunisations in line with the national schedule.</p> <p>Staff members will be encouraged to check they are up-to-date with their MMR vaccinations.</p> <p>Pregnant staff members and those with weak immune systems will be encouraged to contact their GP</p>	<p>Cases are excluded while infectious, which is from four days before the onset of a rash to four days after.</p>

Disease	Symptoms	Considerations	Exclusion period
		immediately for advice if they come into contact with measles.	
Meningitis	Symptoms include fever, severe headaches, photophobia (aversion to light), stiff neck, non-blanching rash, vomiting and drowsiness.	children are encouraged to be up-to-date with their vaccinations. Meningitis is a notifiable disease.	Once a case has received any necessary treatment, they can return to setting once they have recovered.
Meningococcal meningitis and septicaemia (sepsis)	Symptoms include fever, severe headache, photophobia, drowsiness, and a non-blanching rash. Not all symptoms will be present.	Medical advice will be sought immediately. The confidentiality of the case will always be respected. The HPT will be notified of a case of meningococcal disease in the setting. The HPT will be notified if two cases of meningococcal disease occur in the setting within four weeks.	When the case has been treated and recovered, they can return to the setting. Exclusion is not necessary for household or close contacts unless they have symptoms suggestive of meningococcal infection.
Methicillin resistant staphylococcus aureus (MRSA)	Symptoms are rare but include skin infections and boils.	All infected wounds will be covered.	No exclusion is required.
Mpox (monkeypox)	Symptoms are rare but begin within 5 to 21 days after a close physical contact with someone who has mpox infection and may include flu like	The case will be encouraged to consult their GP.	Exclude until the rash has scabbed, all the scabs have fallen off and a fresh layer of skin has formed underneath.

Disease	Symptoms	Considerations	Exclusion period
	symptoms, fever, low energy, swollen glands, general body aches.	Any close contacts will be advised to contact their local HPT for advice.	
Mumps	Symptoms include a raised temperature, swelling and tenderness of salivary glands, headaches, joint pain and general malaise. Mumps may also cause swelling of the testicles.	The case will be encouraged to consult their GP. Parents are encouraged to immunise their children against mumps.	Cases can return to setting five days after the onset of swelling if they feel able to do so.
Norovirus	Symptoms include nausea, diarrhoea, and vomiting. It is known as the 'winter vomiting bug' and the most common cause of gastroenteritis.	The HPT will be contacted if there a higher than previously experience and/or rapidly increasing number of child, staff and beneficiary absences due to diarrhoea and vomiting.	Exclusion until 48 hours after symptoms have stopped and they are well enough to return.
Panton-Valentine Leukocidin Staphylococcus aureus (PVL-SA)	Symptoms can include recurrent boils, skin abscesses and cellulitis.	The HPT will contacted if there are two or more cases.	Exclusion is not necessary unless cases have a lesion or wound that cannot be covered. Cases should not visit gyms or swimming pools until wounds have healed.
Respiratory infections, including coronavirus	Symptoms can be caused by several respiratory infections including the common cold, coronavirus (COVID-19), flu, and respiratory syncytial virus (RSV).	Cases with symptoms will be encouraged to cover their mouth and nose with a tissue when coughing and sneezing, and to wash their hands afterwards.	Cases with mild symptoms, e.g. a runny nose and/or sore throat, can continue to attend if they are otherwise well.

Disease	Symptoms	Considerations	Exclusion period
	Symptoms can be wide-ranging, including a runny nose, high temperature, cough and sore throat, and loss or change in sense of smell or taste.	The DfE helpline and/or the local HPT will be contacted if an outbreak occurs or there is evidence of severe disease, e.g. hospital admission.	Cases who are unwell and have a high temperature should remain at home until they no longer have a high temperature. It is not recommended that children and young people are tested for coronavirus unless directed to by a health professional. Cases aged 18 years and under with a positive test result should stay at home for 3 days after the day they took their test.
Ringworm	Symptoms vary depending on the area of the body affected. The main symptom is a rash, which can be scaly, dry, swollen or itchy and may appear red or darker than surrounding skin.	Cases with ringworm of the feet will wear socks and trainers at all times and cover their feet during physical activities. Cases will be advised to seek advice from a GP for recommended treatment.	No exclusion is usually necessary. For infections of the skin and scalp, cases can return to the setting once they have started treatment.
Rotavirus	Symptoms include severe diarrhoea, stomach cramps, vomiting, dehydration and mild fever.	Cases will be sent home if unwell and encouraged to speak to their GP.	Cases will be excluded until 48 hours have passed since symptoms were present.
Rubella (German measles)	Symptoms are usually mild. Symptoms include a rash, swollen lymph glands, sore throat and runny	MMR vaccines are promoted to all beneficiaries.	Cases will be excluded for five days from the appearance of the rash.

Disease	Symptoms	Considerations	Exclusion period
	nose, mild fever, headache, tiredness, conjunctivitis, painful and swollen joints.		
Scabies	Symptoms include tiny pimples and nodules on the skin. Burrows may be present on the wrists, palms, elbows, genitalia and buttocks.	<p>All household contacts and any other very close contacts should have one treatment at the same time as the second treatment of the case.</p> <p>The second treatment must not be missed and should be carried out one week after the first treatment.</p>	Cases will be excluded until after the first treatment has been carried out.
Scarlet Fever and Invasive group A Streptococcal Disease	<p>Scarlet fever is highly infectious. It is usually a mild illness, though severe complications can occur in rare circumstances. It may be confused with measles.</p> <p>Symptoms include:</p> <ul style="list-style-type: none"> - Flu-like symptoms, e.g. a high temperature, swollen glands and an aching body - Sore throat and/or tonsillitis - A rash that feels rough, like sandpaper, i.e. scarlet fever, typically on the chest and stomach - Flushed cheeks - Scabs and sores 	<p>Cases will be encouraged to visit their GP. The HPT will be contacted if:</p> <ul style="list-style-type: none"> - Two or more cases occur within 10 days of each other, and the affected individuals have a link. - There are cases of serious disease which have resulted in overnight stays in hospital. - There are cases of chickenpox and/or influenza co-circulating in the group where a case of scarlet fever has been confirmed. 	Cases are excluded and can return 24 hours after commencing appropriate antibiotic treatment – cases not receiving treatment will be excluded until resolution of symptoms.

Disease	Symptoms	Considerations	Exclusion period
	- A white coating on tongue		
Slapped cheek syndrome, Parvovirus B19, Fifth's Disease	Where symptoms develop, a rose-red rash making the cheeks appear bright red may appear several days after a mild feverish illness. The rash usually peaks after a week and then fades.	Cases will be encouraged to visit their GP. Parents are requested to inform the charity of a diagnosis of slapped cheek syndrome.	Exclusion is not required – cases are not infectious by the time the rash occurs.
Threadworm	Symptoms include itching around the anus or vagina, particularly at night, and worms may be seen in stools or around the bottom.	Cases will be encouraged to visit their pharmacy for advice on treatment.	Exclusion is not required.
Tuberculosis (TB)	Symptoms include cough, loss of appetite, weight loss, fever, sweating (particularly at night), breathlessness and pains in the chest. TB in parts of the body other than the lungs may produce a painful lump or swelling.	Advice will be sought from the HPT before taking any action, and regarding exclusion periods.	Cases with infectious TB can return to the setting after two weeks of treatment if well enough to do so, and as long as they have responded to anti-TB therapy. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, will not be excluded.
Typhoid and Paratyphoid fever	Symptoms include fatigue, fever and constipation. The symptoms of paratyphoid fever include fever, diarrhoea and vomiting.	All cases will be immediately reported to the HPT.	Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.

Disease	Symptoms	Considerations	Exclusion period
			Environmental health officers or the HPT may advise the setting to issue a lengthened exclusion period.
Whooping cough (pertussis)	Symptoms include a heavy cold with a temperature and persistent cough. The cough generally worsens and develops the characteristic 'whoop'. Coughing spasms may be worse at night and may be associated with vomiting.	Cases will be advised to see their GP. Parents are advised to have their children immunised against whooping cough.	Cases will not return to setting until they have had 48 hours of appropriate treatment with antibiotics and feel well enough to do so, or 21 days from the onset of illness if no antibiotic treatment is given. Cases will be allowed to return in the above circumstances, even if they are still coughing.



Appendix B - Infection absence periods

This table details the minimum required period for staff, children and beneficiaries to stay away from the setting following an infection, as recommended by [UK Health Security Agency](https://www.gov.uk/government/organisations/uk-health-security-agency).

*Identifies a notifiable disease. It is a statutory requirement that doctors report these diseases to their local PHE centre.

Infection	Recommended minimum period to stay away from setting	Comments
Athlete's foot	None	Advise cases to visit their local pharmacy or GP for advice and treatment. Individuals should not be barefoot at their setting e.g. in changing areas and should not share towels, socks or shoes with others.
Chicken pox	At least 5 days from onset of rash and until all blisters have crusted over.	Pregnant staff contacts should consult with their GP or midwife.
Cold sores	None	Avoid contact with the sores.
Conjunctivitis	None	Advise cases to visit their local pharmacy or GP for advice and treatment. If an outbreak occurs, consult the HPT.
Respiratory infections including coronavirus (COVID-19)	Cases should not attend if they have a high temperature and are unwell.	Cases with mild symptoms such as runny nose, and headache who are otherwise well can continue to attend.

Infection	Recommended minimum period to stay away from setting	Comments
	Cases who have a positive test result for COVID-19 should not attend the setting for 3 days after the day of the test.	
Diarrhoea and/or vomiting	Whilst symptomatic and 48 hours from the last episode	Contact the HPT if there are a higher than previously experienced and/or rapidly increasing number of absences due to diarrhoea and vomiting. If a particular cause of the diarrhoea and vomiting is identified, there may be additional exclusion advice, for example E. coli STEC and hep A.
Diphtheria*	Exclusion is essential.	Family contacts must be excluded until cleared by the HPT and the HPT must always be consulted.
Flu (influenza)	Until recovered	Report outbreaks to the HPT.
Glandular fever	Exclusion is not necessary, and cases can return to setting as soon as they feel well.	Glandular fever is spread through spit and can be transferred through kissing or by sharing cups or cutlery. Cases will be infectious for up to 7 weeks before symptoms appear.
Hand, foot and mouth	None	Contact the HPT if a large number of children are affected. Exclusion may be considered in some circumstances.

Infection	Recommended minimum period to stay away from setting	Comments
Head lice	None	Treatment recommended only when live lice seen. Exclusion is not normally permitted. In severe, ongoing cases, the LA does have the power to exclude; however, exclusion should not be overused.
Hepatitis A*	Seven days after onset of jaundice or other symptoms	If it is an outbreak, the HPT will advise on control measures.
Hepatitis B*, C* and HIV	None	Not infectious through casual contact. Procedures for bodily fluid spills must be followed.
Impetigo	48 hours after commencing antibiotic treatment, or when lesions are crusted and healed	Antibiotic treatment is recommended to speed healing and reduce the infectious period.
Measles*	Four days from onset of rash and well enough	Preventable by vaccination (MMR). Follow procedures for vulnerable children and pregnant staff.
Meningococcal meningitis*/septicaemia*	Until recovered	Meningitis ACWY and B are preventable by vaccination. The HPT will advise on any action needed.
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. The HPT will advise on any action needed.

Infection	Recommended minimum period to stay away from setting	Comments
Meningitis viral	None	As this is a milder form of meningitis, there is no reason to exclude those who have been in close contact with infected persons. The local HPT should be consulted.
MRSA	None	Good hygiene – in particular, environmental cleaning and handwashing – is important to minimise the spread. The local HPT should be consulted.
Mumps*	Five days after onset of swelling	Preventable by vaccination with two doses of MMR.
Ringworm	Exclusion is not usually required	Treatment is required.
Rubella* (German measles)	Five days from onset of rash	Preventable by two doses of immunisation (MMR). Pregnant staff contacts should seek prompt advice from their GP or midwife.
Scabies	Can return to setting after first treatment	The infected person's household and those who have been in close contact will also require treatment at the same time.
Scarlet Fever* and Invasive group A Streptococcal Disease	24 hours after commencing antibiotic treatment	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered. If two or more cases occur, the HPT should be contacted.
Slapped cheek/Fifth disease/Parvo Virus B19	None (once rash has developed)	Pregnant contacts of case should consult with their GP or midwife.
Threadworms	None	Treatment recommended for the infected person and household contacts.

Infection	Recommended minimum period to stay away from setting	Comments
Tonsillitis	None	There are many causes, but most causes are virus-based and do not require antibiotics.
Tuberculosis (TB)	<p>If pulmonary TB - until at least 2 weeks after the start of effective antibiotic treatment.</p> <p>Exclusion not required for non-pulmonary or latent TB infection.</p>	Only pulmonary (lung) TB is infectious. It requires prolonged close contact to spread. Consult the local HPT before disseminating information to staff and parents. The HPT will organise any necessary contact tracing.
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms.
Whooping cough (pertussis)*	Two days from commencing antibiotic treatment, or 21 days from the onset of illness if no antibiotic treatment is given	Preventable by vaccination. Non-infectious coughing can continue for many weeks after treatment. The HPT will organise any necessary contact tracing.

Appendix C - Diarrhoea and vomiting outbreak action checklist

Action	Action taken?		Comments
	Yes	No	
A 48-hour exclusion rule has been enforced for ill children, staff and beneficiaries.			
Individuals with symptoms have been kept in an area away from communal areas where they can be observed until family member or parent collects them.			
Liquid soap and paper hand towels are available at all hand wash basins.			
Enhanced cleaning is undertaken twice daily as a minimum, and an appropriate disinfectant is used.			
Advice has been given on the cleaning of vomit, e.g. steam cleaning carpets and furniture and machine hot washing of soft furnishings.			
Appropriate disposable personal protective equipment (PPE) is available.			
Appropriate waste disposal systems are in place for removing infectious waste.			
Staff wearing appropriate PPE when dealing with spills, which will be removed and disposed of quickly.			
Hard toys and equipment are cleaned and disinfected on a daily basis, and their use is limited and rotated.			
The use of soft toys, water and sand play, and cookery activities have been suspended.			

Infected linen is segregated, and dissolvable laundry bags are used where possible.			
Visitors are restricted, and essential visitors are informed of the outbreak and advised on hand washing.			
New attendees joining the affected group or session are delayed from joining.			
The health protection team (HPT) has been informed of any infected food handlers.			
Staff work in dedicated areas and food handling is restricted where possible.			
All staff (including volunteers) are asked if they are unwell and excluded for 48 hours if unwell.			
Trays of fruit/snacks covered until point of serving. Snacks served in individual bowls handed directly to children and beneficiaries.			
Drink bottles clearly labelled with names when required.			
The HPT is informed of any planned events at the setting.			

Date:	
Completed by (print name):	
Signed:	

Please hand this form into main reception where it will be stored securely.